

AD-A185 802

EVALUATION STUDIES OF THE DEN-2/S-1 VACCINE(U) PUERTO
RICO UNIV SAN JUAN E KRAISELBURD NOV 82
DAMD17-80-C-0060

1/1


UNCLASSIFIED

F/G 6/5

NL



END
DATE
FILMED
1-84
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A135 802

12

AD _____

EVALUATION STUDIES OF THE DEN-2/S-1 VACCINE

Annual and Final Report

Edmundo Kraiselburd, Ph.D.

November 1982

Supported by

US Army Medical Research and Development Command
Fort Detrick, Frederick, Maryland 21701

Contract No. DAMD17-80-C-0060

University of Puerto Rico
Medical Sciences Campus, GPO Box 5067
San Juan, Puerto Rico 00936

DOD Distribution Statement

Approved for public release; distribution unlimited

DTIC
ELECTE
DEC 14 1983
E

The findings in this report are not to be construed as
an official Department of the Army position unless so
designated by other authorized documents.

DTIC FILE COPY

83 12 13 173

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
	AD-A135802	
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
EVALUATION STUDIES OF THE DEN-2/S-1 VACCINE		Annual (Aug 1981 - Nov 1982) Final (1 April 80-3 Nov 1982)
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)		8. CONTRACT OR GRANT NUMBER(s)
Edmundo Kraiselburd, Ph.D.		DAMD17-80-C-0060
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
University of Puerto Rico Medical Sciences Campus San Juan, Puerto Rico 00936		62770A.3M162770A871.AA.047
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
US Army Medical Research and Development Command Fort Detrick Frederick, Maryland 21701		November 1982
		13. NUMBER OF PAGES
		7
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)		
Approved for public release, distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
DEN-2 and -4 viruses dose rhesus monkeys seroconversion		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
<p>The purpose of this study was to determine how much dengue 2 and dengue 4 viruses were required to infect rhesus monkeys. To this effect, groups of five dengue antibody free animals were inoculated subcutaneously with serial 10 fold dilutions of dengue 2 (PR159) and dengue 4 (H241) viruses. Virus titrations were carried out simultaneously by direct plaque assay in LLC-MK₂ cells, by mosquito inoculation using both <u>Aedes aegypti</u> and <u>Toxorhynchites amboiensis</u> and in the monkeys. Blood samples for serology were taken at 30</p>		

Enclosure 4

20. Abstract (continuation)

and 60 days post infection. Neutralization tests revealed that 9.5 mosquito infectious dose₅₀ (MID)₅₀ of dengue 2 and 22 MID₅₀ of dengue 4 were required to infect 50% of the monkeys. The data suggest that in recipients challenges with wild type virus, a minimum of 100 MID₅₀'s should be used as the virus challenge dose. (Kraiselburd, E. and Gubler, D., Manuscript in Preparation)

Accomplishments: A group of 31 tuberculin-negative juvenile rhesus monkeys were pretested for neutralizing antibodies to dengue-2 and 4 serotypes. All animals were found to be free of dengue virus antibody. All monkeys were housed at the Caribbean Primate Research Center (Sabana Seca) in individual steel squeeze cages in a mosquito-proof area, 15 days before the pretest bleeding was performed. Groups of 5 animals were inoculated with serial dilutions of either dengue-2 (PR159) or dengue-4 (H241) virus preparations.

Dengue-2 virus (geometric mean titer 2.8×10^4 pfu/ml; 3.0×10^6 MID₅₀/ml) was serially diluted to $1/5 \times 10^{-5}$, $1/5 \times 10^{-6}$, $1/5 \times 10^{-7}$. Group 1 monkeys were s.c. inoculated with 0.5 ml of $1/5 \times 10^{-5}$ dilution (5.6×10^{-2} pfu/ml; 6.0 MID₅₀/ml). Group 2 and group 3 monkeys were s.c. inoculated with 0.5 ml of $1/5 \times 10^{-6}$ dilution (5.6×10^{-3} pfu/ml; 6.0×10^{-1} MID₅₀/ml) and $1/5 \times 10^{-7}$ dilution (5.6×10^{-4} pfu/ml; 6.0×10^{-2} MID₅₀/ml), respectively (see Table II). Dengue-4 virus (geometric mean titer 7.2×10^2 pfu/ml; 1.5×10^6 MID₅₀/ml) was diluted to $1/5 \times 10^{-4}$, $1/5 \times 10^{-5}$ and $1/5 \times 10^{-6}$. Groups 4, 5 and 6 monkeys were s.c. inoculated with 0.5 ml of $1/5 \times 10^{-5}$, $1/5 \times 10^{-6}$ and $1/5 \times 10^{-7}$ dilutions of dengue-4 respectively (see table I). One monkey was used as uninoculated control. Animal inoculation was performed on April 13, 1982.

MID₅₀ (mosquito infection dose 50) were determined using both Aedes Aegypti and Toxorhynchitis mosquitoes by Dr. Gubler (San Juan Laboratories, CDC, Puerto Rico) before and after animal inoculation. MID₅₀ titers reported are geometric means of three independent MID₅₀ determinations.

Animals were bled on post infection days 30 and 60. Seroconversion to the respective dengue virus type was determined by N test performed at 35°C for 30 minutes. Two monkeys (B766 and B773) died on post infection day 30.

An autopsy revealed that the animals died of suffocation secondary to Ketamine sedation. Histopathological review of the tissues confirmed that the animals died of complications of recovery from sedation.

Only one group 1 monkey seroconverted. The remaining dengue-2 infected monkeys did not mount an antibody response to the diluted virus inoculum. Only 2 group 4 monkeys and only 1 group 5 monkeys mounted a positive antibody response to dengue-4 virus.

The minimum dose of dengue virus required for animal seroconversion could not be determined by this experiment (see Quarterly Report, June 30, 1982). To determine this virus dose, lower virus dilutions were used in the experiments described below.

On infection day (June 15, 1982), dengue-2 and dengue-4 viruses were titered again. The titers obtained with the newly defrosted virus samples were 2.5×10^4 pfu/ml (dengue-2) and 800 pfu/ml (dengue-4).

Since on April 13, the virus titers obtained were 3.2×10^4 pfu/ml (dengue-2) and 6.5×10^2 pfu/ml (dengue-4), geometric mean titers (GMT) were calculated for both virus preparations. GMT of 2.8×10^4 pfu/ml and 720 pfu/ml were obtained from dengue-2 and dengue-4, respectively. GMT's were also obtained from three independent MID_{50} determinations (performed by Dr. Gubler, CDC, San Juan). The MID_{50} titers obtained were 3.0×10^6 MID_{50} /ml (dengue-2) and 1.5×10^6 MID_{50} /ml (dengue-4). These GMT were used for calculating monkey ID_{50} from data presented on tables I and II.

On June 15, 1982 groups 2 and 3 monkeys (previously inoculated with diluted dengue-2) were s.c. infected with 0.5 ml of 1.44 pfu/ml and 0.144 pfu/ml of dengue-4, respectively. Groups 6 and 5 monkeys (previously inoculated with diluted dengue-4) were s.c. infected with 0.5 mls of 5.6 pfu/ml and 0.56 pfu/ml of dengue-2 respectively. Infected monkeys were bled on post infection days 30 and 60 and their N antibody titers determined. Results from these experiments are presented in tables I and II.

TABLE I

Immune Response of Rhesus Monkeys to a serially diluted dengue-4 (H24) Virus Preparation.

Dengue 4 Virus Dose		Seroconversion	Geometric N Titers	
pfu	MID ₅₀		30d	60d
7.2×10^{-1}	3.5×10^3	5/5	46	> 160
7.2×10^{-2}	3.5×10^2	4/5	60	> 160
7.2×10^{-3}	3.5×10^1	2/5	49	160
7.2×10^{-4}	3.5	1/5	< 10	160
7.2×10^{-5}	3.5×10^{-1}	0/5	< 10	< 10

Conclusion: For dengue 4, 22 mosquito ID₅₀ (10^{-2} pfu) are equivalent to 1 monkey ID₅₀

- 3 -



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

TABLE II

Immune Response of Rhesus Monkeys to a Serially Diluted Dengue-2 (PR159) Virus Preparation

Dengue 2 Virus Dose		Seroconversion	Geometric N Titers	
pfu	MID50		30d	60d
2.8	3×10^2	5/5	>160	>160
2.8×10^{-1}	3×10^1	4/5	>160	>160
2.8×10^{-2}	3	1/5	>10	>640
2.8×10^{-3}	3×10^{-1}	0/5	<10	<10
2.8×10^{-4}	3×10^{-2}	0/5	<10	<10

Conclusion: For dengue 2, 9.5 mosquito $1D_{50}$ (13×10^{-2} pfu) are equivalent to 1 monkey $1D_{50}$.

Problems: Viruses received from WRAIR were supposed to have 100x higher titers than the one obtained in our laboratory. Therefore, the viruses may have lost titer after shipment from WRAIR to Puerto Rico. This may be of importance in future experiments.

FOREWORD

In conducting the research described in this report, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (DHEW Publication No. (NIH) 78-23, Revised 1978).

DISTRIBUTION LIST

12 copies

Director
Walter Reed Army Institute of Research
Walter Reed Army Medical Center
ATTN: SGRD-UWZ-C
Washington, DC 20012

4 copies

Commander
US Army Medical Research and Development Command
ATTN: SGRD-RMS
Fort Detrick, Frederick, MD 21701

12 copies

Defense Technical Information Center (DTIC)
ATTN: DTIC-DDA
Cameron Station
Alexandria, VA 22314

1 copy

Dean
School of Medicine
Uniformed Services University
of the Health Sciences
4301 Jones Bridge Road
Bethesda, MD 20014

1 copy

Commandant
Academy of Health Sciences, US Army
ATTN: AHS-CDM
Fort Sam Houston, TX 78234